

Exercises, Summer school on the Hitchin fibration

Szilárd Szabó

26–30 July 2010, Bonn

Exercise 1. Let $z = x_1 + ix_2$ be the standard holomorphic coordinate on an open set $U \subset \mathbf{C}$, and set

$$\nabla = d + A_1 dx_1 + A_2 dx_2$$

and

$$\Phi = \frac{1}{2}(\phi_1 - i\phi_2)dz \in \Omega^{1,0}(C, \text{ad}(P)^{\mathbf{C}}).$$

Show that the Self-Duality Equations are equivalent to

$$F_{\nabla} + [\Phi, \Phi^*] = 0 \tag{RSD}$$

$$\nabla^{0,1}\Phi = 0, \tag{CSD}$$

where ∇ acts on endomorphisms by the adjoint action.

Exercise 2. Set $\mathcal{G} = \Omega^0(C, \text{Ad}(P))$ and $\mathcal{G}^{\mathbf{C}} = \Omega^0(C, \text{Ad}(P)^{\mathbf{C}})$. Show that for any $g \in \mathcal{G}^{\mathbf{C}}$, if (∇, Φ) is a solution of (CSD) then so is $g \cdot (\nabla, \Phi)$. Show that for any $g \in \mathcal{G}$, if (∇, Φ) is moreover a solution of (RSD) then so is $g \cdot (\nabla, \Phi)$.

Exercise 3. Let $(V, \langle \cdot, \cdot \rangle)$ be a finite-dimensional Hermitian vector space and G a finite dimensional compact connected Lie group. Recall that V carries a natural Kähler-structure defined by the Riemannian metric

$$g(\mathbf{v}, \mathbf{w}) = \Re \langle \mathbf{v}, \mathbf{w} \rangle$$

and the symplectic form

$$\omega(\mathbf{v}, \mathbf{w}) = g(i\mathbf{v}, \mathbf{w}) = \Im \langle \mathbf{v}, \mathbf{w} \rangle$$

for any $\mathbf{v}, \mathbf{w} \in V$. Assume the compactification $G^{\mathbf{C}}$ of G acts on V smoothly by linear transformations, and assume that the action restricted to G preserves the Hermitian structure too. Define the map $\mu : V \rightarrow \mathfrak{g}^{\vee}$ by

$$\mu(\mathbf{v})(\xi) = \frac{i}{2} \langle \xi \cdot \mathbf{v}, \mathbf{v} \rangle,$$

for any $\xi \in \mathfrak{g}$ and $\mathbf{v} \in V$, where $\xi \cdot \mathbf{v}$ stands for the infinitesimal action of \mathfrak{g} on V . Show that μ is a moment map of the restricted action of G .

Exercise 4. With the notations of Exercise 3, let now $\chi : G \rightarrow \text{U}(1)$ be any character of G , and call the induced complex character $\chi^{\mathbf{C}} : G^{\mathbf{C}} \rightarrow \mathbf{C}^*$. Denote by $d\chi : \mathfrak{g} \rightarrow i\mathbf{R}$ the derivative of χ ; in other words, we have $i(d\chi) \in \mathfrak{g}^{\vee}$. The action of $G^{\mathbf{C}}$ on V and the character $\chi^{\mathbf{C}}$ give rise to the action of $G^{\mathbf{C}}$ on $V \times \mathbf{C}$ defined by the formula

$$g \cdot (\mathbf{v}, z) = (g \cdot \mathbf{v}, \chi^{\mathbf{C}}(g)^{-1}z).$$

We say that $\mathbf{v} \in V$ is χ -stable if for all $z \in \mathbf{C}^*$ the orbit of (\mathbf{v}, z) under this action is closed in $V \times \mathbf{C}$; denote by $V^{\chi-s}$ the set of χ -stable points \mathbf{v} of V . A generalisation of the Theorem seen in class states that there exists a one-to-one correspondence between $\mu^{-1}(i(d\chi))/G$ and $V^{\chi-s}/G^{\mathbf{C}}$.

1. Apply this result to the standard action of $U(1)$ on \mathbf{C}^{n+1} and the character $\chi_l : U(1) \rightarrow U(1)$ given by $g \mapsto g^l$ for a fixed $l \in \mathbf{Z}$. What are the χ_l -stable points, what is the moment map, and what are the quotients on the two sides of the correspondence?
2. Let $0 < k < n$ be integers. Apply the above result to the case where $V = \text{Hom}_{\mathbf{C}}(\mathbf{C}^k, \mathbf{C}^n)$, $G = U(k)$ acting by $g \cdot \varphi = \varphi \circ g^{-1}$ for any $\varphi \in V$ and $g \in G$, and $\chi = \det^{-1} : G \rightarrow U(1)$.

Exercise 5. Let $n \in \mathbf{N}$, $V = \text{End}(\mathbf{C}^n)$ and $G = U(n)$ act on it by conjugation:

$$g \cdot \varphi = g \circ \varphi \circ g^{-1}.$$

Find the moment map μ of this action, the set of stable points of V and compare the two quotients.

Exercise 6. (Atiyah-Bott) Let $G = U(n)$, P be a G -bundle on C and \mathcal{A} denote the affine space of unitary connections ∇ on P . Recall that ∇ is uniquely determined by its $(0,1)$ -part $\nabla^{0,1}$, therefore the tangent space to \mathcal{A} at ∇ can be identified to $\Omega^{0,1}(C, \text{ad}(P)^{\mathbf{C}})$. Let the Kähler structure on $\mathcal{A} \times \Omega^{1,0}(C, \text{ad}(P)^{\mathbf{C}})$ be given by the complex structure $(\psi, \phi) \mapsto (i\psi, i\phi)$ and the Hermitian metric

$$\langle (\psi_1, \phi_1), (\psi_2, \phi_2) \rangle = 2i \int_C \text{tr}(\psi_2^* \wedge \psi_1 + \phi_1 \wedge \phi_2^*).$$

Moreover, let $\mathcal{G} = \Omega^0(C, \text{Ad}(P))$ act on $\mathcal{A} \times \Omega^{1,0}(C, \text{ad}(P)^{\mathbf{C}})$ by conjugation. Observe that the Lie-algebra $\text{Lie}(\mathcal{G})$ of \mathcal{G} can be naturally identified with $\Omega^0(C, \text{ad}(P))$, and its dual with $\Omega^2(C, \text{ad}(P))$ by the Killing form and integration over C . Show that the moment map of the action of \mathcal{G} on $\mathcal{A} \times \Omega^{1,0}(C, \text{ad}(P)^{\mathbf{C}})$ is then

$$\begin{aligned} \mu : \mathcal{A} \times \Omega^{1,0}(C, \text{ad}(P)^{\mathbf{C}}) &\rightarrow \Omega^2(C, \text{ad}(P)) \\ (\nabla, \Phi) &\mapsto F_{\nabla} + [\Phi, \Phi^*]. \end{aligned}$$

Exercise 7. Consider the 4-th power $(\mathbf{CP}^1)^4$ of \mathbf{CP}^1 , and let Δ be the diagonal composed of quadruples in which not all points are distinct. Let the group $G^{\mathbf{C}} = \text{Sl}_2(\mathbf{C})$ act on $X = (\mathbf{CP}^1)^4 \setminus \Delta$ diagonally by the standard action. Determine the set of stable points, the moment map of the action restricted to $G = U(2)$ and the various quotient spaces.